

Strong Lensing Probabilities in a Cosmological Model with a Running Primordial Power Spectrum

Tong-Jie Zhang

*Department of Astronomy, Beijing Normal University, Beijing 100875, P.R.China;
tjzhang@bnu.edu.cn*

Zhi-Liang Yang

*Department of Astronomy, Beijing Normal University, Beijing 100875, P.R.China;
zlyang@bnu.edu.cn*

Xiang-Tao He

*Department of Astronomy, Beijing Normal University, Beijing 100875, P.R.China;
xthe@bnu.edu.cn*

ABSTRACT

The combination of the first-year Wilkinson Microwave Anisotropy Probe (WMAP) data with other finer scale cosmic microwave background (CMB) experiments (CBI and ACBAR) and two structure formation measurements (2dFGRS and Lyman α forest) suggest a Λ CDM cosmological model with a running spectral power index of primordial density fluctuations. Motivated by this new result on the index of primordial power spectrum, we present the first study on the predicted lensing probabilities of image separation in a spatially flat Λ CDM model with a running spectral index (RSI- Λ CDM model). It is shown that the RSI- Λ CDM model suppress the predicted lensing probabilities on small splitting angles of less than about $4''$ compared with that of standard power-law Λ CDM (PL- Λ CDM) model.

Subject headings: cosmology:theory—dark matter—galaxies:halos—gravitational lensing—early universe—large-scale structure

PACS numbers:98.80.Es,98.62.Sb,95.35.+d,98.80.Cq

1. Introduction

Mapping the mass distribution of matter in the universe has been a major challenge for modern observational cosmology. One of the direct procedures to weigh matter in the

universe is measuring its deflection of light by gravity. The statistics of gravitational lensing can provide us with a very powerful probe of the mass distribution of the Universe. It is well known that the Jodrell-Bank VLA Astrometric Survey (JVAS) and the Cosmic Lens All-Sky Survey (CLASS) (Browne & Myers 2000; Myers et al. 2003; Browne et al. 2003) have provided us with observations of strong lensing probabilities for small image separations ranging from $0.3''$ to $3''$. By comparing predicted lensing probabilities with observations, we can examine the mass distributions of dark matter halos, in particular, their inner density slopes (Huterer & Ma 2004; Ricotti 2003).

Based on the Cold Dark Matter (CDM) model, which has become the standard theory of cosmic structure formation, the lensing probabilities strongly depend on the density profiles of CDM halos. The lensing model is usually described by a singular isothermal sphere (SIS), the Navarro-Frenk-White (NFW) model (Navarro et al. 1996, 1997), or generalized NFW (GNFW) density profiles of dark halos (Zhao 1996). Li & Ostriker (2002) employed a semi-analytical approach to analyze the gravitational lensing of remote quasars by foreground dark halos and checked the plausibility of various lensing models. They found that no model can completely explain the current observations: the SIS models predict too many lenses with large splitting angles, while the NFW models predict too few small splitting angles. They therefore further developed a two-population halo model for lensing: small mass halos with a steep inner density slope and large mass halos with a shallow inner density slope, concluding that a combination of SIS and NFW halo models can reproduce the current observations reasonably well. Unlike previous work that directly models the density profiles of dark matter halos semi-analytically, Zhang (2004) generalized the density profiles of dark matter halos from high-resolution N-body simulations by means of generalized Navarro-Frenk-White (GNFW) models of three populations with slopes, α , of about -1.5, -1.3 and -1.1 for galaxies, groups and clusters, respectively. He presented the calculations of lensing probabilities using these GNFW profiles for three populations in various spatially flat cosmological models with a cosmological constant Λ . He showed that the compound model of density profiles does not match well with the lensing probabilities derived from the combined data of JVAS/CLASS. Recently, Metcalf (2004) compared predictions on small scale structure for the Λ CDM model by numerical simulations with observed flux ratios, and found that the disagreements between monochromatic flux ratios and simple lens models can be explained without any substructure in the dark matter halos of primary lenses. However, spectroscopic lensing observations of Q2237+0305 require more small mass dark halos than that expected in the Λ CDM model.

In particular, a power spectrum of primordial fluctuation, $P_p(k)$, should be assumed in advance in the calculation of lensing probabilities. Inflationary models predict a approximately scale-invariant power spectra for primordial density (scalar metric) fluctuation, $P_p(k) \propto k^n$ with index $n = 1$ (Guth & Pi 1982; Bardeen et al. 1983). The combination of

the first-year Wilkinson Microwave Anisotropy Probe (WMAP) data with other finer scale cosmic background (CMB) experiments (Cosmic Background Imager [CBI], Arcminute Cosmology Bolometer Array Receiver [ACBAR]) and two observations of large-scale structure (the Anglo-Australian Telescope Two-Degree Field Galaxy Redshift Survey [2dFGRS] and Lyman α forest) favor a Λ CDM cosmological model with a running index of the primordial power spectrum (RSI- Λ CDM), while the WMAP data alone still suggest a best-fit standard power-law Λ CDM model with the spectral index of $n \approx 1$ (PL- Λ CDM) (Spergel et al. 2003; Peiris et al. 2003). However, there still exist the intriguing discrepancies between theoretical predictions and observations on both the largest and smallest scales. While the emergence of a running spectral index may improve problems on small scales, there remain a possible discrepancy on the largest angular scales. It is particularly noted that the running spectral index model suppress significantly the power amplitude of fluctuations on small scales (Spergel et al. 2003; Yoshida et al. 2003). This imply a reduction of the amount of substructure within galactic halos (Zentner & Bullock 2002). Yoshida et al. (2003) studied early structure formation in a RSI- Λ CDM universe using high-resolution cosmological N-body/hydrodynamic simulations. They showed that the reduced small-scale power in the RSI- Λ CDM model causes a considerable delay in the formation epoch of low-mass minihalos ($\sim 10^6 M_\odot$) compared with the PL- Λ CDM model, although early structure still forms hierarchically in the RSI- Λ CDM model. Thus the running index probably affect the strong lensing process of distant sources by intervening dark matter halos because the lensing probabilities strongly depend on the abundance of dark halos formed in the evolution of the universe. In this letter, we will present the first calculation of lensing probabilities in a RSI- Λ CDM model to explore the effect of running spectral index of primordial fluctuation on strong lensing probabilities. We adopt a spatially flat Λ CDM cosmological model through this letter.

The reminder of this paper is organized as follows. We describe strong lensing probabilities and mass function of dark halos in Section 2 and 3 respectively. The cross-sections for producing multiple images are described in Section 4. Our results is shown in Section 5, while a conclusion and discussion are given in Section 6.

2. Strong Lensing Probability

The lensing probability with image separation larger than $\Delta\theta$ is given by Schneider et al. (1992)

$$P(> \Delta\theta) = \int p(z_s) dz_s \int_0^{z_s} \frac{dD_p(z)}{dz} dz \int_0^\infty \bar{n}(M, z) \sigma(M, z) dM. \quad (1)$$

In this expression, $D_p(z) = c/H_0 \int_0^z dz/(1+z)E(z)$ is the proper distance from the observer to the lens at redshift z where the expansion rate of the universe $E(z) = \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda}$ in a spatially flat cosmological model, and $p(z_s)$ is the redshift distribution of distant sources. The physical number density $\bar{n}(M, z)$ of virialized dark halos of masses between M and $M + dM$ is expressed as $\bar{n}(M, z) = n(M, z)(1+z)^3$ and $\sigma(M, z)$ is the cross-section defined in the lens plane for forming multiple images.

3. Mass Function of Dark Halos

In the standard hierarchical theory of structure formation, the comoving number density of virialized dark halos per unit mass M at redshift z can be given by the Press and Schechter (PS) formula (Press & Schechter 1974): $n(M, z) = dN/dM = \rho_0 f(M, z)/M$ where ρ_0 is the mean mass density of the universe today and, instead of PS formula in this letter, the mass function $f(M, z)$ takes the form of an empirical fit from high-resolution simulation (Jenkins et al. 2001)

$$f(M, z) = \frac{0.301}{M} \frac{d \ln \Delta^{-1}(M, z)}{d \ln M} \exp(-|\ln \Delta^{-1}(M, z) + 0.64|^{3.88}). \quad (2)$$

Here, $\Delta(M, z) = \Delta(M)D(z)$ and $D(z) = e(\Omega(z))/e(\Omega_m)(1+z)$ is the linear growth function of density perturbation (Carroll et al. 1992), in which $e(x) = 2.5x/(1/70 + 209x/140 - x^2/140 + x^{4/7})$ and $\Omega(z) = \Omega_m(1+z)^3/E^2(z)$. The present variance of the fluctuations within a sphere containing a mass M can be expressed as $\Delta^2(M) = \frac{1}{2\pi^2} \int_0^\infty P(k)W^2(kr_M)k^2 dk$, where $W(kr_M) = 3[\sin(kr_M)/(kr_M)^3 - \cos(kr_M)/(kr_M)^2]$ is the Top-hat window function in Fourier space and $r_M = (3M/4\pi\rho_0)^{1/3}$. The power spectrum of CDM density fluctuations is $P(k) = P_p(k)T^2(k)$ where the matter transfer function $T(k)$ is given by Eisenstein & Hu (1999), and $P_p(k)$ is the primordial power spectrum of density fluctuation. The scale-invariant primordial power spectrum in the PL- Λ CDM model is given by $P_p(k) = Ak^{n_s}$ with index $n_s=1$ and that in the RSI- Λ CDM model is assumed to be $P_p(k) = P(k_0)(k/k_0)^{n_s(k)}$, where the index $n_s(k)$ is a function of length scale

$$n_s(k) = n_s(k_0) + \frac{1}{2} \frac{dn_s(k)}{d \ln k} \ln \left(\frac{k}{k_0} \right). \quad (3)$$

The pivot scale $k_0=0.05 \text{ h Mpc}^{-1}$, $n_s(k_0)=0.93$, and $dn_s/d \ln k=-0.03$ are the best-fit values to the combination data of the recent CMB experiments and two other large-scale structure observations (Spergel et al. 2003). For both PL- Λ CDM and RSI- Λ CDM models, the amplitude of primordial power spectrum, A and $P(k_0)$, are normalized to $\sigma_8 = \Delta(r_M = 8h^{-1}\text{Mpc})$, which is the rms mass fluctuations when present universe is smoothed using a window function on a scale of $8h^{-1}\text{Mpc}$.

4. The Cross-sections for Producing Multiple Images

The cross-section for producing multiple images relies on the density profile of dark matter halos. Based on previous work (Li & Ostriker 2002; Zhang 2004), we model dark halos as an SIS for $M < M_{c1}$ and NFW profile for $M > M_{c1}$, respectively, where $M_{c1} \sim 10^{13} h^{-1} M_{\odot}$ corresponds to the cooling mass scale (Porciani & Madau 2000; Kochanek & White 2001).

4.1. The Cross-section for SISs

The mass density for a lens of SIS is $\rho(r) = \sigma_v^2 / 2\pi G r^2$ where σ_v is the velocity dispersion (Schneider et al. 1992). Thus the surface mass density of the SIS satisfies $\Sigma(\xi) = \sigma_v^2 / 2G\xi$ where $\xi \equiv |\vec{\xi}|$, $\vec{\xi}$ is the position vector in the lens plane. By defining the length scales in the lens plane and the source plane as $\xi_0 = 4\pi(\sigma_v/c)^2 d_L^A d_{LS}^A / d_S^A$, $\eta_0 = \xi_0 d_S^A / d_L^A$ respectively, we can simplify the lensing equation and obtain the image separation of SIS lensing

$$\Delta\theta = 1.27' \left(\frac{d_{LS}^A}{d_S^A} \right) M_{15}^{2/3} E(z)^{2/3}, \quad (4)$$

where $M_{15} = M / (10^{15} h^{-1} M_{\odot})$ is dimensionless halo mass. Here d_S^A and d_L^A are the angular diameter distances from the observer to the source and to the lens object respectively, while d_{LS}^A is the same quantity but from the lens to the source object. Therefore, the cross-section defined in the lens plane for forming two images by an SIS lens with splitting angle $\Delta\theta > \Delta\theta_0$ can be expressed as $\sigma(M, z) = \pi \xi_0^2 \vartheta(\Delta\theta - \Delta\theta_0)$ where ϑ is a step function.

4.2. The Cross-section for NFW Models

The GNFW density profile can be expressed in the form $\rho(r) = \rho_s r_s^3 / r^\alpha (r + r_s)^{3-\alpha}$ (Zhao 1996) where $0 < \alpha < 3$. The GNFW density profile reduces to the case of NFW if $\alpha = 1$. The mass of a dark halo within r_{200} can be defined as $M = 4\pi \int_0^{r_{200}} \rho r^2 dr = 4\pi \rho_s r_s^3 f(c_1)$, and r_{200} is the radius of a sphere around a dark halo within which the average mass density is 200 times the critical mass density of the universe. The function $f(c_1) = \int_0^{c_1} x^2 dx / x^\alpha (1+x)^{3-\alpha}$ and $c_1 = r_{200}/r_s$ is the concentration parameter and takes the form (Oguri et al. 2001, 2002; Oguri 2004)

$$c_1(M_{15}, z) = c_{\text{norm}} \frac{2-\alpha}{1+z} [10M_{15}]^{-0.13}, \quad (5)$$

where z is the redshift of halo, $c_{\text{norm}} = 8$ (Bullock et al. 2001). So ρ_s and r_s can be related to mass M_{15} and redshift z by

$$\rho_s = \rho_{c_0} E^2(z) \frac{200}{3} \frac{c_1^3}{f(c_1(M_{15}, z))}, \quad r_s = \frac{1.626}{c_1} \frac{M_{15}^{1/3}}{E^{2/3}(z)} h^{-1} \text{Mpc}, \quad (6)$$

where ρ_{c_0} is the critical mass density of the universe today. The lensing equation for the GNFW profile is given by $y = x - \mu_s g(x)/x$; $\vec{\xi} = \vec{x} r_s$ and $\vec{\eta} = \vec{y} r_s d_S^A/d_L^A$ are the position vectors in the lens plane and the source plane respectively, $g(x) \equiv \int_0^x u du \int_0^\infty (u^2 + z^2)^{-\alpha/2} \left[(u^2 + z^2)^{1/2} + 1 \right]^{-3}$ and

$$\mu_s \equiv \frac{4\rho_s r_s}{\Sigma_{\text{cr}}} = 0.002 \left(\frac{\rho_s}{\rho_{c_0}} \right) \left(\frac{r_s}{1h^{-1}\text{Mpc}} \right) \left(\frac{d_R^A}{c/H_0} \right), \quad (7)$$

where μ_s is a parameter on which the efficiency of producing multiple images is strongly dependent. Here $\Sigma_{\text{cr}} = \frac{c^2}{4\pi G} \frac{d_S^A}{d_L^A d_{LS}^A}$ is the critical surface mass density, and $d_R^A = d_L^A d_{LS}^A/d_S^A$. The lensing equation curves are symmetrical with respect to the origin. Multiple images can be formed when $|y| \leq y_{\text{cr}}$, where y_{cr} is the maximum value of y when $x < 0$ or the minimum value for $x > 0$. Generally speaking, there exist three images for $|y| < y_{\text{cr}}$. We will just consider the outermost two images stretched by the splitting angle $\Delta\theta$ when more than two images are formed. Therefore, we can write the cross-section as $\sigma(M, z) \approx \pi y_{\text{cr}}^2 r_s^2 \vartheta(\Delta\theta - \Delta\theta_0)$ with $\Delta\theta > \Delta\theta_0$ in the lens plane for multiple images produced by a GNFW lens at z . Here splitting angle $\Delta\theta$ is given by $\Delta\theta = r_s \Delta x/d_L^A \approx 2x_0 r_s/d_L^A$ and x_0 is the positive root of the lensing equation $y(x) = 0$.

5. Numerical Results

In this letter, we assume spatially flat Λ CDM models characterized by the matter density parameter Ω_m , vacuum energy density parameter Ω_Λ . For both PL- Λ CDM and RSI- Λ CDM models, we take cosmological parameters to be the new result from the WMAP: Hubble constant $h = 0.71$, $\Omega_m = 0.27$, $\sigma_8 = 0.84$ (Bennett et al. 2003; Spergel et al. 2003) and $M_{\text{cl}} = 1.5 \times 10^{13} h^{-1} M_\odot$. As mentioned above, the lensing probability depend strongly on the abundance of virialized dark halos. The mass function of dark halos directly involve the calculation of primordial power of density fluctuation. According to Eq.(1), it is clear that the effect of running spectral index on mass function of halos cause the difference of lensing probabilities between the two models.

Although the redshift distribution of quasars in the JVAS/CLASS survey is still poorly known, the prediction of Dunlop & Peacock (1990) model and the CLASS lensing sub-sample redshift measurements suggest that the redshift distribution for CLASS unlensed sources can be modelled by a Gaussian distribution with mean redshift $\langle z_s \rangle = 1.27$ (Marlow et al. 2000) and dispersion $\sigma_z = 0.95$ (Chae et al. 2002; Browne et al. 2003; Myers et al. 2003). Thus in

this letter we adopt the Gaussian redshift distribution of quasars with the mean redshift $\langle z_s \rangle = 1.27$ and dispersion $\sigma_z = 0.95$. In the definition of cross-section for forming two images, we just consider the criterion $\Delta\theta$, and neglect another one q_r that is the flux density (brightness) ratio of multiple images (Schneider et al. 1992). In order to investigate the effect of central black holes or bulges on lensing probability, Chen (2003a,b) introduced q_r into the calculation for lensing cross-section. Due to the existence of central black holes or galactic bulges, y_{cr} becomes extremely large when $|x|$ approaches zero. Thus y_{cr} can be determined by the consideration of q_r together with $\Delta\theta$. However for GFW halo models in the absence of central black holes or galactic bulges, the lensing equation curves are so smooth that we do not need to define cross-section by q_r . As for SISs, it is necessary to introduce q_r because both the predicted lensing probabilities and the determination of the cosmic equation of state ω are quite sensitive to q_r (Chen 2004a,b). Our objective in this letter is to examine the effect of running spectral index on the predicted lensing probabilities, so for explicit we neglect this selection criterion q_r for forming multiple images. In such cases, we calculate lensing probabilities with image separations greater than $\Delta\theta$ according to the compound model of SIS and NFW halo profile in both PL- Λ CDM and RSI- Λ CDM models. Our numerical results are shown in Fig.1 together with the observational one from JVAS/CLASS. As we expect, there is slight difference between the two models on large image separations, while this difference enlarges with the decrease of the splitting angle. More specifically, the RSI- Λ CDM model can reduce the predicted lensing probabilities on small splitting angle of less than about $4''$ compared with that of PL- Λ CDM model.

6. Conclusions and Discussion

Motivated by the new result on the index of primordial power spectrum from a combination of WMAP data with other finer scale CMB experiments and other large-scale structure observations, we present the first study on the predicted lensing probability in a Λ CDM model with a running spectral index. In a popular cosmological model from the new fit results mentioned above: $h = 0.71$, $\Omega_m = 0.27$, $\sigma_8 = 0.84$, we calculate lensing probabilities with image separations greater than $\Delta\theta$ according to the compound model of SIS and NFW halo profile in both PL- Λ CDM and RSI- Λ CDM models. From the analysis above, we can see that the running spectral index mainly affect the predicted lensing probability on small image separation. It is well known that structures in the universe forms hierarchically in standard CDM models. Yoshida et al. (2003) found that although this hierarchical formation mechanism do not work well in RSI- Λ CDM model compared with that in PL- Λ CDM model and it also is not clear that the PS theory can be used in RSI- Λ CDM model, the mass function measured by high-resolution cosmological N-body/hydrodynamic simulations

overall match the PS mass function for both RSI- Λ CDM and PL- Λ CDM model.

In addition, because the running spectral index model predicts a significant lower power of density fluctuation on small scales than the standard PL- Λ CDM model (Spergel et al. 2003; Yoshida et al. 2003), it should also attract considerable attention in studies on weak lensing by large-scale structure (Ishak et al. 2003), especially on skewness (Pen et al. 2003; Zhang et al. 2003) which characterizes the non-Gaussian property of κ field in the nonlinear regime.

We are very grateful to the anonymous referee for constructive suggestion. T.J.Zhang would like to thank Ue-Li Pen, Peng-Jie Zhang, Xiang-Ping Wu, Bo Qin and CITA for their hospitality during his visits to the Canadian Institute for Theoretical Astrophysics(CITA), University of Toronto and the cosmology groups of the National Astronomical Observatories of P.R.China. This work was supported by the National Science Foundation of China (grants No.10473002 and 10273003).

REFERENCES

- Bardeen, J. M., Steinhardt, P. J., & Turner, M. S. 1983, *Phys. Rev. D*, 28, 679
- Bennett, C. L., Halpern, M., Hinshaw, G., Jarosik, N., Kogut, A., Limon, M., Meyer, S. S., Page, L., Spergel, D. N., Tucker, G. S., Wollack, E., Wright, E. L., Barnes, C., Greason, M. R., Hill, R. S., Komatsu, E., Nolte, M. R., Odegard, N., Peiris, H. V., Verde, L., & Weiland, J. L. 2003, *ApJS*, 148, 1
- Browne, I. W. A. & Myers, S. T. 2000, in *IAU Symposium*
- Browne, I. W. A., Wilkinson, P. N., Jackson, N. J. F., Myers, S. T., Fassnacht, C. D., Koopmans, L. V. E., Marlow, D. R., Norbury, M., Rusin, D., Sykes, C. M., Biggs, A. D., Blandford, R. D., de Bruyn, A. G., Chae, K.-H., Helbig, P., King, L. J., McKean, J. P., Pearson, T. J., Phillips, P. M., Readhead, A. C. S., Xanthopoulos, E., & York, T. 2003, *MNRAS*, 341, 13
- Bullock, J. S., Kolatt, T. S., Sigad, Y., Somerville, R. S., Kravtsov, A. V., Klypin, A. A., Primack, J. R., & Dekel, A. 2001, *MNRAS*, 321, 559
- Carroll, S. M., Press, W. H., & Turner, E. L. 1992, *ARA&A*, 30, 499
- Chae, K.-H., Biggs, A. D., Blandford, R. D., Browne, I. W., de Bruyn, A. G., Fassnacht, C. D., Helbig, P., Jackson, N. J., King, L. J., Koopmans, L. V., Mao, S., Marlow, D. R., McKean, J. P., Myers, S. T., Norbury, M., Pearson, T. J., Phillips, P. M.,




Fig. 1.— The lensing probabilities for image separations greater than $\Delta\theta$. The solid-line histogram represents observed lensing probabilities from JVAS/CLASS. The solid line is the lensing probabilities for running spectral index Λ CDM model, while the dashed line is that for power law Λ CDM model.

- Readhead, A. C., Rusin, D., Sykes, C. M., Wilkinson, P. N., Xanthopoulos, E., & York, T. 2002, *Physical Review Letters*, 89, 151301
- Chen, D. 2003a, *ApJ*, 587, L55
- . 2003b, *A&A*, 397, 415
- . 2004a, *Chinese Journal of Astronomy and Astrophysics*, 4, 118
- Chen, D.-M. 2004b, *A&A*, 418, 387
- Dunlop, J. S. & Peacock, J. A. 1990, *MNRAS*, 247, 19
- Eisenstein, D. J. & Hu, W. 1999, *ApJ*, 511, 5
- Guth, A. H. & Pi, S.-Y. 1982, *Physical Review Letters*, 49, 1110
- Huterer, D. & Ma, C. 2004, *ApJ*, 600, L7
- Ishak, M., Hirata, C. M., McDonald, P., & Seljak, U. 2003, *ArXiv Astrophysics e-prints:astro-ph/0308446*
- Jenkins, A., Frenk, C. S., White, S. D. M., Colberg, J. M., Cole, S., Evrard, A. E., Couchman, H. M. P., & Yoshida, N. 2001, *MNRAS*, 321, 372
- Kochanek, C. S. & White, M. 2001, *ApJ*, 559, 531
- Li, L. & Ostriker, J. P. 2002, *ApJ*, 566, 652
- Marlow, D. R., Rusin, D., Jackson, N., Wilkinson, P. N., Browne, I. W. A., & Koopmans, L. 2000, *AJ*, 119, 2629
- Metcalf, R. B. 2004, *ArXiv Astrophysics e-prints:astro-ph/0407298*
- Myers, S. T., Jackson, N. J., Browne, I. W. A., de Bruyn, A. G., Pearson, T. J., Readhead, A. C. S., Wilkinson, P. N., Biggs, A. D., Blandford, R. D., Fassnacht, C. D., Koopmans, L. V. E., Marlow, D. R., McKean, J. P., Norbury, M. A., Phillips, P. M., Rusin, D., Shepherd, M. C., & Sykes, C. M. 2003, *MNRAS*, 341, 1
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1996, *ApJ*, 462, 563
- . 1997, *ApJ*, 490, 493
- Oguri, M. 2004, *ArXiv Astrophysics e-prints:astro-ph/0408573*

- Oguri, M., Taruya, A., & Suto, Y. 2001, *ApJ*, 559, 572
- Oguri, M., Taruya, A., Suto, Y., & Turner, E. L. 2002, *ApJ*, 568, 488
- Peiris, H. V., Komatsu, E., Verde, L., Spergel, D. N., Bennett, C. L., Halpern, M., Hinshaw, G., Jarosik, N., Kogut, A., Limon, M., Meyer, S. S., Page, L., Tucker, G. S., Wollack, E., & Wright, E. L. 2003, *ApJS*, 148, 213
- Pen, U., Zhang, T., van Waerbeke, L., Mellier, Y., Zhang, P., & Dubinski, J. 2003, *ApJ*, 592, 664
- Porciani, C. & Madau, P. 2000, *ApJ*, 532, 679
- Press, W. H. & Schechter, P. 1974, *ApJ*, 187, 425
- Ricotti, M. 2003, *MNRAS*, 344, 1237
- Schneider, P., Ehlers, J., & Falco, E. E. 1992, *Gravitational Lenses* (Gravitational Lenses, XIV, 560 pp. 112 figs.. Springer-Verlag Berlin Heidelberg New York. Also *Astronomy and Astrophysics Library*)
- Spergel, D. N., Verde, L., Peiris, H. V., Komatsu, E., Nolta, M. R., Bennett, C. L., Halpern, M., Hinshaw, G., Jarosik, N., Kogut, A., Limon, M., Meyer, S. S., Page, L., Tucker, G. S., Weiland, J. L., Wollack, E., & Wright, E. L. 2003, *ApJS*, 148, 175
- Yoshida, N., Sokasian, A., Hernquist, L., & Springel, V. 2003, *ApJ*, 598, 73
- Zentner, A. R. & Bullock, J. S. 2002, *Phys. Rev. D*, 66, 43003
- Zhang, T. 2004, *ApJ*, 602, L5
- Zhang, T., Pen, U., Zhang, P., & Dubinski, J. 2003, *ApJ*, 598, 818
- Zhao, H. 1996, *MNRAS*, 278, 488